

THE ROLE OF ACTIVITY FOR RHYTHM DISRUPTION IN NIGHT SHIFT WORKERS

Melissa Alves Braga de Oliveira, Cláudia Ávila Moraes, Rosa Maria Levandovski, Camila Morelato de Souza, Maria Paz Loayza Hidalgo

Introduction: Night duty shifts have been identified as a good model of human circadian rhythm disruption. As a result, they may suffer from several health problems associated with disturbed physiological rhythms. **Objective:** This study aimed to compare activity-rest, light exposure and temperature rhythms among regular workers and night shift workers who engage in high- or lowintensity activity. **Methods:** This was a cross-sectional study of 20 women with a mean age of 44 ± 3.45 years. Comparisons were conducted between regular day workers and night shift workers engaging in low-intensity and high-intensity work. Actigraphy was employed to assess rhythmic variables for 7 consecutive days. Circadian rhythms were calculated by fitting the data to a sinusoidal curve capturing a 24-hour period and obtaining its amplitude, acrophase and mesor. Rayleigh test for circular statistics was used. Student's independent t-test and one-way analysis of variance (ANOVA) with Tukey's post-hoc tests were used to assess significant differences among groups. **Results:** Regular workers had a lower temperature amplitude and higher activity amplitude than shift workers. Additionally, shift workers with high activity levels show a phase advance in activity and temperature when compared with shift workers with low-intensity work and regular workers. There were no light exposure differences among groups. **Conclusion:** Our results suggest that light may not be the only factor associated with biological rhythm alterations observed in night shift workers. Rather, the amount of activity performed appears to affect their biological rhythms. Therefore, when considering shift work as a model for circadian rhythm disruption, it is essential to properly qualify and quantify activity intensity during duty.