

ORIGINAL ARTICLE

Open Access



# Validation of reference genes for normalization of gene expression by qRT-PCR in a resveratrol-producing entophytic fungus (*Alternaria* sp. MG1)

Jin-xin Che<sup>1</sup>, Jun-ling Shi<sup>2\*</sup>, Yao Lu<sup>2</sup> and Yan-lin Liu<sup>1\*</sup>

## Abstract

*Alternaria* sp. MG1, an endophytic fungus isolated from *Vitis vinifera*, can independently produce resveratrol, indicating that this species contains the key genes for resveratrol biosynthesis. Identification of these key genes is essential to understand the resveratrol biosynthesis pathway in this strain, which is currently unknown in microorganisms. qRT-PCR is an efficient and widely used method to identify the key genes related to unknown pathways at the level of gene expression. Verification of stable reference genes in this strain is essential for qRT-PCR data normalization, although results have been reported for other *Alternaria* sp. strains. In this study, nine candidate reference genes including *TUBA*, *EF1*, *EF2*, *UBC*, *UFD*, *RPS5*, *RPS24*, *ACTB* and *18S* were evaluated for expression stability in a diverse set of six samples representing different growth periods. We compared cell culture conditions and an optimized condition for resveratrol production. The comparison of the results was performed using four statistical softwares. A combination of *TUBA* and *EF1* was found to be suitable for normalization of *Alternaria* sp. MG1 in different developmental stages, and *18S* was found to be the least stable. The reference genes verified in this study will facilitate further research to explore gene expression and molecular mechanisms as well as the improvement of secondary metabolite yields in *Alternaria* sp. MG1. To our knowledge, this is the first validation of reference genes in *Alternaria* with the capability to produce resveratrol. Additionally, these results provide useful guidelines for the selection of reference genes in other *Alternaria* species.

**Keywords:** qRT-PCR, Reference genes,  $\alpha$ -Tubulin, Elongation factor 1, *Alternaria* sp.

## Introduction

Many important bioactive compounds are widely used in medical services and health care (Khan 2016; Larsen and Matchkov 2016; Morata et al. 2015). Many of these compounds are either microbial metabolites or their semi-synthetic derivatives (Golinska et al. 2015; Sessitsch et al. 2013; Stepniewska and Kuzniar 2013). In the microbial population, endophytes are a large group which may contain millions of different species, but only a minority of them have

been studied (Sessitsch et al. 2013). Several endophytic fungi (*Alternaria* sp.) were identified previously that are capable of independent resveratrol production (Shi et al. 2012). Although fundamental physiological research has been performed (Zhang et al. 2013a, b), the metabolic pathways and cellular processes remain to be elucidated.

Gene expression profiling is an informative technique to investigate biological systems (Li et al. 2015). The method of qRT-PCR (quantitative real time PCR) can measure gene expression across different sample populations (Derveaux et al. 2010; Wong and Medrano 2005). However, there are many factors that can influence the accuracy of the results such as the quality and quantity of mRNA templates or amplification efficiency. Generally, normalizing expression of the target genes to one or

\*Correspondence: sjshi2004@nwpu.edu.cn; yanlinliu@aliyun.com

<sup>1</sup> College of Food Science and Engineering, Northwest A & F University, 28 Xinong Road, Yangling 712100, Shaanxi, People's Republic of China

<sup>2</sup> Key Laboratory for Space Biosciences and Space Biotechnology, School of Life Sciences, Northwestern Polytechnical University, Xi'an 710072, Shaanxi, People's Republic of China

several reference genes provide an efficient way to reduce these effects and increase the relevance of the results (Huggett et al. 2005; Marabita et al. 2016). However, the use of inappropriate reference genes that change expression levels under different conditions can cause interpretation errors. Thus, the choice of appropriate reference genes for normalization is a prerequisite for qRT-PCR assay.

In recent years, validation of reliable reference genes before their use for normalization has been performed for many species, such as *Talaromyces marneffei* (Dankai et al. 2015), *Staphylococcus aureus* (Sihto et al. 2014), *Beauveria bassiana* (Zhou et al. 2012), *Oenococcus oeni* (Sumbly et al. 2012) and others. Commonly used reference genes for these fungi include the genes encoding the 18S ribosomal RNA (*18S*), ubiquitin fusion degradation protein (*UFD*), ribosomal protein (*RPS*), elongation factor (*EF*),  $\beta$ -actin (*ACTB*),  $\alpha$ -tubulin (*TUBA*), ubiquitin-conjugating enzyme (*UBC*), and glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) (Kozera and Rapacz 2013). The previously reported reference genes for normalizing qRT-PCR data in fungi, especially in *Alternaria* sp. are  $\beta$ -tubulin (*TUB*) for *A. alternata* (Baez-Flores et al. 2011; Buzina et al. 2008) and *A. brassicicola* (Sellam et al. 2006), *benA* for *A. alternata* (Saha et al. 2012), 18S for *A. infectoria* (Fernandes et al. 2014), and elongation factor 1 (*EF1*) for *A. brassicicola* (Cho et al. 2014). Genes that show stable expression under many conditions may differ in microorganisms due to different organization structures and when different genes are expressed. Therefore, it is necessary to identify reliable reference genes in *Alternaria* sp. MG1 for use in qRT-PCR assay.

The aim of this study was to identify the most stable reference genes in *Alternaria* sp. under different growth conditions and resveratrol production conditions. Genes that show relatively similar levels of expression under all conditions could serve as reference genes that would be appropriate for comparison in the qRT-PCR assay of genes whose expression may vary during changes in metabolism or during resveratrol biosynthesis. Several software applications were used for analysis of candidate reference genes. These programs allowed evaluation of appropriate reference genes under given experimental conditions using statistical methods, such as Bestkeeper (Pfaffl et al. 2004), geNorm (Vandesompele et al. 2002), and Normfinder (Andersen et al. 2004).

## Materials and methods

### Microorganism

*Alternaria* sp. MG1 (code: CCTCC M 2011348), a strain previously isolated from the cob of Merlot grape (Shi et al. 2012), was used in the study. It was maintained at the China Center for Type Culture Collection (Wuhan,

China). For preparation of *Alternaria* sp. MG1 cells, the strain was inoculated into a 250-mL flask containing 100 mL PDB (liquid potato-dextrose broth, potato 200 g with 20 g glucose in 1000 mL tap-water). The cultivation was carried out at 28 °C and 120 rpm in a rotary shaker. According to the growth curve analysis (Additional file 1: Figure S1), the cells were collected at points throughout lag phase, logarithmic growth phase, and stationary phase after a cultivation of 2, 3, 4, 5 and 6 days by centrifugation at 1136 $\times$ g for 10 min at 4 °C (HC-3018R, Anhui USTC Zonkia Scientific Instruments Co., Ltd., Anhui, China). Next, the collected cells were washed twice with sterile water and immediately stored in liquid nitrogen until further analysis. The resting cells were collected using the method reported by Zhang et al. (2013a) as follows: after cultivation for 4 days, the rinsed cells were resuspended in 0.2 mol/L pH 7.0 phosphate buffer containing 0.1 g/L MgSO<sub>4</sub>, 0.2 g/L CaSO<sub>4</sub> and 4 mmol/L phenylalanine for 21 h. After that, cells were washed twice with sterile water and stored for further analysis.

### Total RNA extraction and cDNA synthesis

Extraction of the total RNA from the cells was performed using a Spin Column Fungal total RNA Purification Kit (Sangon Biotech (Shanghai) Co., Ltd., China). Quality and quantity of the RNA extraction were analyzed using a NanoDrop 2000 Spectrophotometer (Thermo Scientific, Waltham, Massachusetts, USA) gel electrophoresis, and Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, California, USA). First-strand cDNA was synthesized using a PrimeScript<sup>TM</sup> RT reagent Kit with gDNA Eraser (Perfect Real Time) in strict accordance with the manufacturer's operation manual (TAKARA Biotechnology (Dalian) CO., LTD., China). The cDNA products were diluted fivefold and stored at -20 °C before further analysis.

### Primer design of reference genes and qRT-PCR amplification conditions

The nine candidate genes (Table 1), *ACTB*, *EF1*, *EF2*, *RPS5*, *RPS24*, *TUBA*, *UBC*, *UFD* and *18S* were selected based on the transcriptome database of *Alternaria* sp. MG1 (available through NCBI, SRA study accession number SRP060338) (Che et al. 2016b). The internal reference genes had highly similar sequences with reported genes from previous studies (DiGuistini et al. 2011; Farajalla and Gulick 2007; Goodwin et al. 2011; Skora et al. 2015). The primer pairs of candidate genes were designed using the software Primer Premier (version 5.00) (<http://www.premierbiosoft.com/primerdesign/index.html>) with an amplicon length ranging from 100 to 300 bp.

The real-time quantitative PCR amplification and analysis were performed in the Bio-Rad iQ<sup>TM</sup>5 Multicolor

**Table 1** Relation of primers for the candidate genes to internal control

Internal gene	Gene name	Primer sequence (5'–3') Forward/reverse	Amplicon length (bp)	Amplification efficiency (%)	Regression coefficient ( $R^2$ )	Accession number at GenBank
<i>ACTB</i>	$\beta$ -Actin	CAAGACGGAAGGCTGG AA/ CACTGCCGAGCGAGAAAT	195	100.4	0.997	GEMY01018051
<i>EF1</i>	Elongation factor 1	CACTGGTTTTGCCTT TTCCT/ TGTGGGCACCGTCAAAGT	186	127.3	0.995	GEMY01015044
<i>EF2</i>	Elongation factor 2	ATAACAGCCTGGAAG ATGC/ CTTTCACCATCCGTCAGTT	207	98.3	0.996	GEMY01001243
<i>RPS5</i>	Ribosomal protein S5	ACACCCATACAAAGAACG/ CCGAGTGCCTTGCTGA	131	104.1	0.985	GEMY01011888
<i>RPS24</i>	Ribosomal protein S24	CCGTCTTGTCGTTCCC/ CGATTGGCGGTTTCTC	133	104.8	0.997	GEMY01015522
<i>TUBA</i>	$\alpha$ -Tubulin	CAAGCGAGTCAGAAGC/ GGTATGTTGGTGAGGGTAT	101	106.9	0.984	GEMY01012167
<i>UBC</i>	Ubiquitin-conjugating enzyme	GGCTCAAGAAACAGGAA/ AGATTACCACCCGAAC	123	100.4	0.984	GEMY01016137
<i>UFD</i>	Ubiquitin fusion degradation protein	TCCTCCTTGCCCTTGA/ CGAATCCGCCTCCTAC	108	123.6	0.996	GEMY01001986
<i>18S</i>	18S ribosomal RNA	TCTTGTTTCCTTGGTGGGT/ GCATTTCGCTGCCTTCT	144	106.2	0.980	JN102357.1

Real-Time PCR Detection System (Bio-Rad Laboratories, Inc., Hercules, California, USA) with the iQ<sup>TM</sup>5 Optical system Software Version 2.1. (<http://www.bio-rad.com/zh-cn/sku/1709753-iq5-optical-system-software?parentCategoryGUID=2>). A total reaction system of 25  $\mu$ L contained SYBR *Premix Ex TaqII* (Tli RNase Plus) (2 $\times$  Conc.), 12.5  $\mu$ L; PCR primer mix (10  $\mu$ M), 2  $\mu$ L; cDNA template, 1  $\mu$ L; and DNase-free water, 9.5  $\mu$ L. The qRT-PCR amplification program was 95  $^{\circ}$ C for 5 min, followed by 40 cycles of 94  $^{\circ}$ C for 30 s, the ideal annealing temperature for each primer pair for 30 s, and 72  $^{\circ}$ C for 1 min, and then 72  $^{\circ}$ C for 10 min. All reactions were conducted in triplicate and melting curve analysis was performed. The correlation coefficients ( $R^2$ ) and slope values of the standard curve and efficiency ( $E$ ) were calculated using the iQ 5 Optical system Software Version 2.1.

To confirm the accuracy of the amplified products, all the PCR products were analyzed by agarose gel electrophoresis using 2% agarose gels in Tris-borate-EDTA (TBE) buffer stained with ethidium bromide.

#### Determination of reference gene expression stability using data analysis software

The transcript abundance of the reference genes was determined by the Ct value. The expression stability of these candidate reference genes were evaluated using the four methods described below. The methods are Excel-based tool—Bestkeeper software (version 1) ([\[bioinformatics.gene-quantification.info/bestkeeper.html\]\(http://bioinformatics.gene-quantification.info/bestkeeper.html\)\) using pair-wise correlations \(Pfaffl et al. 2004\), Genorm software \(version 3.4\) \(<https://genorm.cmgg.be/>\) \(Vandesompele et al. 2002\), NormFinder software \(version 0.953\) \(<http://moma.dk/normfinder-software/>\) \(Andersen et al. 2004\), and the comparative  \$\Delta\$ Ct method \(Pfaffl 2001; Silver et al. 2006\). To comprehensively analyze the stability of these candidate genes, the web tool RefFinder \(<http://fulxie.0fees.us/?type=reference>\) \(Xie et al. 2012\) was used to compare and rank the outcomes of the results using the different analysis programs.](http://</a></p>
</div>
<div data-bbox=)

## Results

### RNA purity and concentration


The mean values of quantity and quality of the RNA samples are shown in Table 2. The concentrations of RNA samples ranged from 324.00 to 1329.56 ng/ $\mu$ L. The mean values of 260/280 were close to 2.00.

### Verification of primer specificity of selected reference genes

A total of nine candidate genes (Table 1) were selected for this study by referring to previous studies (DiGiustini et al. 2011; Farajalla and Gulick 2007; Goodwin et al. 2011; Skora et al. 2015) and the *Alternaria* sp. MG1 transcriptome database. Agarose gel electrophoresis for preliminary PCR and melting curve analysis was performed and the results are shown in Fig. 1 and

**Table 2** The quantity and quality of RNA samples isolated from *Alternaria* sp. MG1 during different growth stages

Sample	2 days	3 days	4 days	5 days	6 days	Resting cell
260/280	2.18	2.16	2.13	1.97	2.09	2.00
260/230	1.61	1.57	1.65	1.41	1.60	1.52
Conc. (ng/ $\mu$ L)	1329.56	762.96	563.76	324.00	336.84	594.04



Additional file 1: Figure S1, respectively. We observed that the lengths of amplified fragment were consistent with the expected size, and no primer dimers were detected except for candidate gene *RPS24*. These results indicated the primers were specific and suitable for reference gene validation.

PCR efficiency analysis was performed to validate the optimal of the reference gene. The regression coefficient ( $R^2$  value) and PCR amplification efficiency were calculated by a standard curve generated using tenfold serial dilutions of pooled cDNA. The PCR amplification efficiency of these candidate genes ranged from 98.3 to 127.3%, and the regression coefficient ( $R^2$  value) of the standard curve ranged from 0.980 to 0.997, well within the acceptable range of qRT-PCR (Table 1).

**Expression profiling of the candidate reference genes**

Six samples were chosen for each candidate reference gene in this study.


The average expression of the candidate genes during different growth stages was investigated by comparison of Ct values and the results are shown as a box-plot (Fig. 2). In the figure, the interquartile values are shown in boxes. The median expression level and the total expression level are shown as a line and whisker, respectively. The expression level of the nine reference genes with Ct value ranged from 14.22 to 25.66. Lower Ct values indicate higher expression level and vice versa. The *18S* gene showed the highest expression level, and the *EF1* was the lowest. There was little difference among the other candidate reference genes with Ct values between

20.24 and 23.38. Genes that showed different ranges of expression ( $Ct_{max} - Ct_{min}$ ) were *ACTB* (1.52), *EF1* (0.91), *EF2* (2.55), *RPS5* (2.19), *RPS24* (3.48), *TUBA* (1.33), *UBC* (2.52), *UFD* (3.13), and *18S* (5.64).

**Stability evaluation of candidate genes using different analysis programs**

To validate the stability of these nine candidate genes, we used four evaluation methods. The Bestkeeper software was employed to validate and rank the stability evaluation of these candidate genes based on the standard deviation (SD) of Ct values and the coefficient of variance (CV) expressed as a percentage of the Ct values. In this approach, the most stable reference gene was identified by the comparison of SD value and CV value of these selected genes. The lowest SD and CV values represent the genes with a highest stability, and vice versa. Here, the descriptive statistics of these nine candidate genes were calculated based on the Ct values, and the statistical outcome is listed in Table 3.

The geometric mean (GM), arithmetic mean (AM), extremum (min and max) value, standard deviation (SD), and coefficient of variation (CV), were calculated. Sorted by SD values, the tested genes were in the order of *EF1* < *TUBA* < *ACTB* < *RPS5* < *UBC* < *EF2* < *RPS24* < *UFD* < *18S*. The overall variation of *EF1* and *TUBA* were lowest with a SD value of 0.25. The CV values of *EF1* and *TUBA* were lower than the others, 0.99 and 1.2%, respectively. Interrelated analysis provided by BestKeeper concluded that the most stable reference gene was *EF1*, and *TUBA* was the second most stable.

**Table 3** CT data of reference genes calculated using Bestkeeper


Genes	<i>ACTB</i>	<i>EF1</i>	<i>EF2</i>	<i>RPS5</i>	<i>RPS24</i>	<i>TUBA</i>	<i>UBC</i>	<i>UFD</i>	<i>18S</i>
Number of sample	6	6	6	6	6	6	6	6	6
GM (CT)	20.35	25.66	21.56	20.72	20.24	21.13	22.7	23.38	16.9
AM (CT)	20.35	25.66	21.57	20.73	20.27	21.13	22.72	23.4	16.98
Min (CT)	19.57	25.19	19.95	19.86	18.19	20.52	21	22.42	14.42
Max (CT)	21.09	26.1	22.5	22.05	21.67	21.85	23.52	25.55	20.06
SD ( $\pm$ CT)	0.41	0.25	0.61	0.49	0.69	0.25	0.6	0.78	1.05
CV (% CT)	2.04	0.99	2.8	2.38	3.42	1.2	2.66	3.35	6.2
Min (x-fold)	-1.71	-1.39	-3.04	-1.82	-4.15	-1.53	-3.25	-1.95	-5.59
Max (x-fold)	1.67	1.36	1.92	2.51	2.69	1.65	1.76	4.5	8.93
SD (x-fold)	1.33	1.19	1.52	1.41	1.62	1.19	1.52	1.72	2.08

Another program, geNorm (Vandesompele et al. 2002), was used to calculate the average expression stability of M value and analyze the stability of the candidate reference genes. The calculated M values of the nine candidate reference genes are plotted in Fig. 3. The most stable expression genes had the lowest M value, and vice versa. Previous studies (Vandesompele et al. 2002; Wu et al. 2014), suggested selection of stable reference genes with M values below the threshold of 1.5. As shown in Fig. 3, all the M values of the tested genes were less than 1.5. In the growth stage and the resting cells, the *EF1* and *TUBA* showed the highest expression stability with the lowest M values (0.025). Sometimes, normalization with a single reference may produce significant errors and the more than one reference genes may be needed in some experiments. However, some


researchers (Kong et al. 2015; Li et al. 2015) have indicated that multiple reference genes could increase instability and experimental complexity. Thus, pairwise variation ( $V_n/V_{n+1}$ ) was selected for assessing the optimal number of reference genes. The pairwise variation V value was calculated using geNorm, and a threshold V value of 0.15 was recommended to identify the number of the additional reference genes (Vandesompele et al. 2002). In Fig. 4, all the V values were below the cutoff value of 0.15. Pairwise variation analysis showed that the  $V_{2/3}$  value was 0.0126, which indicated two reference genes was sufficient for gene expression normalization and the two stability reference genes selected were *EF1* and *TUBA*.

In an alternate approach, we used the Normfinder software program to evaluate these candidate reference





**Fig. 3** Average expression stability values (M) of the nine candidate reference genes as calculated by geNorm



**Fig. 4** Pairwise variation (V) calculated by geNorm to determine the optimal number of reference genes



genes. As a model-based variance estimation approach, Normfinder is used to calculate stability values and evaluate the expression stabilities of the tested genes (Andersen et al. 2004; Maroufi et al. 2010). A lower average expression stability indicated genes that were stably expressed gene. In Fig. 5, the stability value ranking of these candidate reference genes was slightly different from that calculated by geNorm software. However, the most stable reference gene was the same (*TUBA*), followed by *EF1*, *EF2*, *RPS5*, *RPS24*, *ACTB*, *UFD*, *UBC* and *18S*. The comparative  $\Delta C_t$  method was used to assess gene expression stability. The stability results were the same as those calculated using Normfinder software, and these results are shown together in Fig. 5. Again, the most stable reference gene was *TUBA*.

#### Overall ranking order and selection of optimal reference genes

In the separate assessments, the most stable reference gene was the same, but the other genes were ranked differently in the different analyses. Next, we used the web tool (RefFinder) to arrange the comprehensive results by integration of the results of the four assessments to compare these potential reference genes. An overview of the expression stability of the nine candidate genes from different growing stages and different treatment of *Alternaria* sp. MG1 are shown in Fig. 6. The ranking of these candidate reference genes (from most stable to least stable) were *TUBA*, *EF1*, *EF2*, *ACTB*, *RPS5*, *RPS24*, *UBC*, *UFD*, and *18S*.

#### Discussion

As a bioactive polyphenol, resveratrol has a variety of functions, such as preventing or slowing the occurrence of cancer, acting as a powerful antioxidant, and extending life span. Pharmaceutical production and functional food processing present a high demand of resveratrol. To date, resveratrol was provided by extraction from plant materials. This method of production is highly limited by plant growth times and low yields. Many resveratrol-producing *Escherichia coli* or yeasts have been constructed by genetic modification (Conrado et al. 2012; Krivoruchko and Nielsen 2015). However, these processes have low yield and have stability issues during production due to the use of the plant-derived genes. The problem of low yield results from a complicated metabolic pathway and a rate-limited enzyme. In a previous study, resveratrol production was increased



slightly by adding substrates or production in *Alternaria* sp. MG1 resting cell culture (Zhang et al. 2013b). Thus, understanding the expression of the genes involved in resveratrol biosynthesis pathway is the key problem to be solved. As a newly characterized biological resource, *Alternaria* sp. MG1 is able to produce resveratrol without limitation from plant resources (Che et al. 2016a; Shi et al. 2012). Study of this fungus may allow insight into the necessary pathways allowing engineering for increased production.

With several advantages, including the ability to quantify, reproducibility, sensitivity and accuracy, qRT-PCR is a preferred method to use for quantifying the gene expression, and assessing mRNA levels among different samples. Validation of appropriate reference genes is a prerequisite for accurate analysis of gene expression using qRT-PCR (Li et al. 2016). In the past research, the most traditional reference genes used in qRT-PCR assay were genes such as *ACT*, *TUB*, and *18S*. In the past, many studies showed that the expression of these traditional

reference genes was not always stable under all conditions (Dankai et al. 2015; Zhou et al. 2012). The identification of the most stable reference genes in *Alternaria* sp. MG1 has not been achieved until now, although some traditional reference genes have been used for qRT-PCR data normalization in some other *Alternaria* sp. (Dankai et al. 2015; Sihto et al. 2014).

Several methods have been recently used to determine the stability of gene expression and to validate the best reference genes (Tong et al. 2009). However, there is no consensus on the ideal approach that should be used to examine the stability of reference gene expression. The pairwise comparison strategy, accessible through the geNorm software, is a very popular option to verify the expression stability of candidate genes (Yu et al. 2016). However, co-regulated genes may confound the geNorm software and this would lead to an erroneous choice of optimum normalizer pair (Andersen et al. 2004). To investigate whether the potential co-regulated genes affected the outcome of the results, researchers removed one of the co-regulated genes from analysis and reported that co-regulation did not influence the ranking of reference genes by stability (Tong et al. 2009); we similarly found no effect on the ranking by the inclusion of co-regulated genes (Additional file 1: Figure S2). Additionally, the reference genes that belonged to the same functional class were not top-ranked and did not occupy closed positions by geNorm software in previous studies (Exposito-Rodriguez et al. 2008). Similarly, the co-regulated genes were not top-ranked and did not occupy closed positions in this research. As a result, use of these two pairs of co-regulated genes did not affect the final ranking of the reference genes by using geNorm software. Other methods such as NormFinder and BestKeeper, were reported to be less sensitive to co-regulation, and might serve as appropriate statistical applets to further assess the stability for reference gene expression (Huang et al. 2014). In an effort to ensure the accuracy of the reference gene stability ranking and minimize bias introduced by the validation approach, four different statistical approaches,  $\Delta C_t$ , geNorm, NormFinder, and BestKeeper, were used to identify the suitable reference genes for accurate normalization in this study. The overall ranking of these four approaches was integrated using a web-based comprehensive tool (RefFinder) developed to identify the most reliable reference genes by integrating these four evaluation methods (Xie et al. 2012).

Nine traditional reference genes (*TUBA*, *UFD*, *RPS24*, *RPS5*, *UBC*, *EF1*, *EF2*, *ACTB* and *18S*) were selected as candidate reference genes, and these genes were tested during different growth periods and during an optimized condition for resveratrol production. Different statistical algorithms and analytical methods gave different

validation results. The overall ranking of these results was integrated using the RefFinder system. The comprehensive results of this research demonstrated that *TUBA* and *EF1* were the most stable reference genes, *18S* was the least stable gene, and the other candidate reference genes were intermediate among all six sets of experiments. Interestingly, *TUBA* was shown to be a reliable reference gene for *Penicillium expansum* (De Clercq et al. 2016) and *Valsa mali* var. *mali* (*Vmm*) (Yin et al. 2013), and *EF1* was found to be suitable for *Clonostachys rosea* (Sun et al. 2015) and *Tuber melanosporum* (Cesare et al. 2015). However, *TUBA* and *EF1* were unstable and unsuitable for use as reference genes in *Blumeria graminis* (Pennington et al. 2016), *C. rosea* (Sun et al. 2015), and *Pandora neoaphidis* (Chen et al. 2016). Other candidate reference genes showed different stability in different fungi. For example, the most stable reference gene for *Talaromyces marneffeii* was *GAPDH*, followed by *TUBA*, and *ACTB* (Dankai et al. 2015). *ACTB* was identified as the reliable reference gene in *Penicillium echinulatum* (Zampieri et al. 2014).

Additionally, normalization with the combination of more genes resulted in improved accuracy. Previous research indicated the application of individual or combinations of 2, 3, and 4 reference genes would result in different levels of abundance, but qualitatively similar patterns (Hu et al. 2009). How many reference genes should be used is dependent on the purpose of research. One reference gene would be enough to show a rough expression mode of genes, if the reference gene was identified as a stable expressed gene (Cho et al. 2014). Nevertheless, if the research purpose is to compare gene expression levels among different samples or to get an accurate expression level, the more reference genes used, the more accurate the result is. However, other researches have reported that multiple reference genes could increase instability and experimental complexity (Kong et al. 2015; Li et al. 2015). Thus, pairwise variation ( $V_n/V_{n+1}$ ) was selected to assess the optimal number of reference genes. The pairwise variation  $V$  value was calculated using geNorm, and a threshold  $V$  value of 0.15 was recommended to identify the number of the additional reference genes (Sumby et al. 2012; Vandesompele et al. 2002). In this study, pairwise variation analysis (Fig. 4) showed that the  $V_{2/3}$  value was 0.0126, which indicated that two reference genes was sufficient for gene expression normalization and the two stability reference genes selected were *EF1* and *TUBA*.

In conclusion, the results obtained here and in previous studies indicate that validation of reference genes is crucial for accurate normalization of gene expression measurements under different experimental conditions. The reference genes verified in this study will be useful



for future research to explore gene expression, molecular mechanisms, and improvement of secondary metabolite yields in *Alternaria* sp. MG1. To our knowledge, this was the first validation of reliable reference genes in *Alternaria*. The results of this study provide useful guidelines for the selection of reference genes in other *Alternaria* species.

## Additional file

**Additional file 1. Figure S1.** The growth curve of *Alternaria* sp. MG1 in liquid potato-dextrose broth (PDB) at 28 °C and 120 rpm. **Figure S2.** Average expression stability values (M) of reference genes removed one of the coregulated genes as calculated by geNorm.

### Authors' contributions

Planning and designing of study: JXC, JLS, YLL; experimentation: JXC, YL; result analysis: JXC, YL; manuscript drafting: JXC, JLS, YL. All authors contributed in the final approval of manuscript. All authors read and approved the final manuscript.

### Acknowledgements

A special thanks to Cheng-quan Yang and Yao-hua You (College of Horticulture, Northwest A & F University) for their support and help with this research.

### Competing interests

The authors declare that they have no competing interests.

### Ethics approval

This article does not contain any studies with human participants or animals performed by any of the author.

### Funding

This study was funded by the National Natural Science Fund (Grant No. 31471718), the Agriculture Department of China (Grant No. CARS-30), and the Northwestern Polytechnical University (No. 201410699086 and No. 3102014GEKY1010).

Received: 31 October 2016 Accepted: 31 October 2016

Published online: 08 November 2016

## References

- Andersen CL, Jensen JL, Orntoft TF (2004) Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res* 64(15):5245–5250
- Baez-Flores ME, Troncoso-Rojas R, Osuna MAI, Dominguez MR, Pryor B, Tiznado-Hernandez ME (2011) Differentially expressed cDNAs in *Alternaria alternata* treated with 2-propenyl isothiocyanate. *Microbiol Res* 166(7):566–577
- Buzina W, Raggam RB, Paulitsch A, Heiling B, Marth E (2008) Characterization and temperature-dependent quantification of heat shock protein 60 of the immunogenic fungus *Alternaria alternata*. *Med Mycol* 46(6):627–630
- Cesare P, Cesare P, Ragnelli AM, Aimola P, Leonardi M, Bonfigli A, Colafarina S, Poma AM, Miranda M, Pacioni G (2015) Validation of reference genes for quantitative real-time PCR in Perigord black truffle (*Tuber melanosporum*) developmental stages. *Phytochemistry* 116:78–86
- Che J, Shi J, Gao Z, Zhang Y (2016a) A new approach to produce resveratrol by enzymatic bioconversion. *J Microbiol Biotech* 26(8):1348–1357
- Che J, Shi J, Gao Z, Zhang Y (2016b) Transcriptome analysis reveals the genetic basis of the resveratrol biosynthesis pathway in an endophytic fungus (*Alternaria* sp. MG1) isolated from *vitis vinifera*. *Front Microbiol* 7(1257):12
- Chen C, Xie TN, Ye SD, Jensen AB, Eilenberg J (2016) Selection of reference genes for expression analysis in the entomophthorean fungus *Pandora neoaphidis*. *Braz J Microbiol* 47(1):259–265
- Cho Y, Ohm RA, Devappa R, Lee HB, Grigoriev IV, Kim BY, Ahn JS (2014) Transcriptional responses of the *Bdttf1*-deletion mutant to the phytoalexin brassinin in the necrotrophic fungus *Alternaria brassicicola*. *Molecules* 19(8):10717–10732
- Conrado RJ, Wu GC, Boock JT, Xu HS, Chen SY, Lebar T, Turnsek J, Tomsic N, Avbelj M, Gaber R, Koprivnjak T, Mori J, Glavnik V, Vovk I, Bencina M, Hodnik V, Anderluh G, Dueber JE, Jerala R, DeLisa MP (2012) DNA-guided assembly of biosynthetic pathways promotes improved catalytic efficiency. *Nucleic Acids Res* 40(4):1879–1889
- Dankai W, Pongpom M, Vanittanakom N (2015) Validation of reference genes for real-time quantitative RT-PCR studies in *Talaromyces marneffeii*. *J Microbiol Methods* 118:42–50
- De Clercq N, Vlaemynck G, Van Pamel E, Van Weyenberg S, Herman L, Devlieghere F, De Meulenaer B, Van Coillie E (2016) Isoeopoxydon dehydrogenase (*idh*) gene expression in relation to patulin production by *Penicillium expansum* under different temperature and atmosphere. *Int J Food Microbiol* 220:50–57
- Derveaux S, Vandesompele J, Hellemans J (2010) How to do successful gene expression analysis using real-time PCR. *Methods* 50(4):227–230
- DiGiustini S, Wang Y, Liao NY, Taylor G, Tanguay P, Feau N, Henrissat B, Chan SK, Hesse-Orce U, Alamouti SM, Tsui CKM, Docking RT, Levasseur A, Haridas S, Robertson G, Birol I, Holt RA, Marra MA, Hamelin RC, Hirst M, Jones SJM, Bohlmann J, Breuil C (2011) Genome and transcriptome analyses of the mountain pine beetle-fungal symbiont *Grosmannia clavigera*, a lodge-pole pine pathogen. *Proc Natl Acad Sci USA* 108(6):2504–2509
- Exposito-Rodriguez M, Borges AA, Borges-Perez A, Perez JA (2008) Selection of internal control genes for quantitative real-time RT-PCR studies during tomato development process. *BMC Plant Biol* 8:12
- Farajalla MR, Gulick PJ (2007) The  $\alpha$ -tubulin gene family in wheat (*Triticum aestivum* L.) and differential gene expression during cold acclimation. *Genome* 50(5):502–510
- Fernandes C, Anjos J, Walker LA, Silva BMA, Cortes L, Mota M, Munro CA, Gow NAR, Goncalves T (2014) Modulation of *Alternaria infectoria* cell wall chitin and glucan synthesis by cell wall synthase inhibitors. *Antimicrob Agents Chemother* 58(5):2894–2904
- Golinska P, Wypij M, Agarkar G, Rathod D, Dahm H, Rai M (2015) Endophytic actinobacteria of medicinal plants: diversity and bioactivity. *Anton Leeuw Int J G* 108(2):267–289
- Goodwin SB, Ben M'Barek S, Dhillon B, Wittenberg AHJ, Crane CF, Hane JK, Foster AJ, Van der Lee TAJ, Grimwood J, Aerts A, Antoniw J, Bailey A, Bluhm B, Bowler J, Bristow J, van der Burgt A, Canto-Canche B, Churchill ACL, Conde-Ferraz L, Cools HJ, Coutinho PM, Csukai M, Dehal P, De Wit P, Donzelli B, van de Geest HC, Van Ham R, Hammond-Kosack KE, Henrissat B, Kilian A, Kobayashi AK, Koopmann E, Kourmpetis Y, Kuzniar A, Lindquist E, Lombard V, Maliepaard C, Martins N, Mehrabi R, Nap JPH, Ponomarenko A, Rudd JJ, Salamov A, Schmutz J, Schouten HJ, Shapiro H, Stergiopoulos I, Torriani SFF, Tu H, de Vries RP, Waalwijk C, Ware SB, Wiebenga A, Zwieters LH, Oliver RP, Grigoriev IV, Kema GHJ (2011) Finished genome of the fungal wheat pathogen *Mycosphaerella graminicola* reveals dispensable structure, chromosome plasticity, and stealth pathogenesis. *PLoS Genet* 7(6):17
- Hu RB, Fan CM, Li HY, Zhang QZ, Fu YF (2009) Evaluation of putative reference genes for gene expression normalization in soybean by quantitative real-time RT-PCR. *BMC Mol Biol* 10:12
- Huang LK, Yan HD, Jiang XM, Zhang Y, Zhang XQ, Ji Y, Zeng B, Xu B, Yin GH, Lee S, Yan YH, Ma X, Peng Y (2014) Reference gene selection for quantitative real-time reverse-transcriptase PCR in orchardgrass subjected to various abiotic stresses. *Gene* 553(2):158–165
- Huggett J, Dheda K, Bustin S, Zumla A (2005) Real-time RT-PCR normalisation; strategies and considerations. *Genes Immun* 6(4):279–284
- Khan MZ (2016) A possible significant role of zinc and GPR39 zinc sensing receptor in Alzheimer disease and epilepsy. *Biomed Pharmacother* 79:263–272
- Kong FN, Cao M, Sun PP, Liu WX, Mao YX (2015) Selection of reference genes for gene expression normalization in *Pyropia yezoensis* using quantitative real-time PCR. *J Appl Phycol* 27(2):1003–1010
- Kozera B, Rapacz M (2013) Reference genes in real-time PCR. *J Appl Genet* 54(4):391–406

- Krivoruchko A, Nielsen J (2015) Production of natural products through metabolic engineering of *Saccharomyces cerevisiae*. *Curr Opin Biotech* 35:7–15
- Larsen MK, Matchkov VV (2016) Hypertension and physical exercise: the role of oxidative stress. *Med Lith* 52(1):19–27
- Li XY, Cheng JY, Zhang J, da Silva JAT, Wang CX, Sun HM (2015) Validation of reference genes for accurate normalization of gene expression in *Lilium davidii* var. *unicolor* for real time quantitative PCR. *PLoS ONE* 10(10):17
- Li MY, Wang F, Jiang Q, Wang GL, Tan C, Xiong AS (2016) Validation and comparison of reference genes for qPCR normalization of *Celery* (*Apium graveolens*) at different development stages. *Front Plant Sci* 7:12
- Marabita F, de Candia P, Torri A, Tegner J, Abrignani S, Rossi RL (2016) Normalization of circulating microRNA expression data obtained by quantitative real-time RT-PCR. *Brief Bioinform* 17(2):204–212
- Maroufi A, Van Bockstaele E, De Loose M (2010) Validation of reference genes for gene expression analysis in chicory (*Cichorium intybus*) using quantitative real-time PCR. *BMC Mol Biol* 11:12
- Morata L, Mensa J, Soriano A (2015) New antibiotics against gram-positives: present and future indications. *Curr Opin Pharmacol* 24:45–51
- Pennington HG, Li LH, Spanu PD (2016) Identification and selection of normalization controls for quantitative transcript analysis in *Blumeria graminis*. *Mol Plant Pathol* 17(4):625–633
- Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 29(9):6
- Pfaffl MW, Tichopad A, Prgomet C, Neuvians TP (2004) Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper—excel-based tool using pair-wise correlations. *Biotechnol Lett* 26(6):509–515
- Saha D, Fetzner R, Burkhardt B, Podlech J, Metzler M, Dang H, Lawrence C, Fischer R (2012) Identification of a polyketide synthase required for alternariol (AOH) and alternariol-9-methyl ether (AME) formation in *Alternaria alternata*. *PLoS ONE* 7(7):14
- Sellam A, Poupard P, Simoneau P (2006) Molecular cloning of AbGst1 encoding a glutathione transferase differentially expressed during exposure of *Alternaria brassicicola* to isothiocyanates. *FEMS Microbiol Lett* 258(2):241–249
- Sessitsch A, Kuffner M, Kidd P, Vangronsveld J, Wenzel WW, Fallmann K, Puschenreiter M (2013) The role of plant-associated bacteria in the mobilization and phytoextraction of trace elements in contaminated soils. *Soil Biol Biochem* 60:182–194
- Shi JL, Zeng Q, Liu YL, Pan ZL (2012) *Alternaria* sp. MG1, a resveratrol-producing fungus: isolation, identification, and optimal cultivation conditions for resveratrol production. *Appl Microbiol Biotechnol* 95(2):369–379
- Sihto HM, Tasara T, Stephan R, Johler S (2014) Validation of reference genes for normalization of qPCR mRNA expression levels in *Staphylococcus aureus* exposed to osmotic and lactic acid stress conditions encountered during food production and preservation. *FEMS Microbiol Lett* 356(1):134–140
- Silver N, Best S, Jiang J, Thein SL (2006) Selection of housekeeping genes for gene expression studies in human reticulocytes using real-time PCR. *BMC Mol Biol* 7:9
- Skora J, Otlewska A, Gutarowska B, Leszczynska J, Majak I, Stepień L (2015) Production of the allergenic protein Alt a 1 by *Alternaria* isolates from working environments. *Int J Environ Res Public Health* 12(2):2164–2183
- Stepniewska Z, Kuzniar A (2013) Endophytic microorganisms-promising applications in bioremediation of greenhouse gases. *Appl Microbiol Biotechnol* 97(22):9589–9596
- Sumbly KM, Grbin PR, Jiranek V (2012) Validation of the use of multiple internal control genes, and the application of real-time quantitative PCR, to study esterase gene expression in *Oenococcus oeni*. *Appl Microbiol Biotechnol* 96(4):1039–1047
- Sun ZB, Li SD, Sun MH (2015) Selection of reliable reference genes for gene expression studies in *Clonostachys rosea* 67-1 under sclerotial induction. *J Microbiol Methods* 114:62–65
- Tong ZG, Gao ZH, Wang F, Zhou J, Zhang Z (2009) Selection of reliable reference genes for gene expression studies in peach using real-time PCR. *BMC Mol Biol* 10:13
- Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 3(7):12
- Wong ML, Medrano JF (2005) Real-time PCR for mRNA quantitation. *Biotechniques* 39(1):75–85
- Wu JX, Su SY, Fu LL, Zhang YJ, Chai LJ, Yi HL (2014) Selection of reliable reference genes for gene expression studies using quantitative real-time PCR in navel orange fruit development and pummelo floral organs. *Sci Hortic* 176:180–188
- Xie FL, Xiao P, Chen DL, Xu L, Zhang BH (2012) miRDeepFinder: a miRNA analysis tool for deep sequencing of plant small RNAs. *Plant Mol Biol* 80(1):75–84
- Yin ZY, Ke XW, Huang DX, Gao XN, Voegelé RT, Kang ZS, Huang LL (2013) Validation of reference genes for gene expression analysis in *Valsa mali* var. *mali* using real-time quantitative PCR. *World J Microb Biot* 29(9):1563–1571
- Yu SH, Yang P, Sun T, Qi Q, Wang XQ, Xu DL, Chen XM (2016) Identification and evaluation of reference genes in the Chinese white wax scale insect *Ericerus pela*. *SpringerPlus* 5:8
- Zampieri D, Nora LC, Basso V, Camassola M, Dillon AJ (2014) Validation of reference genes in *Penicillium echinulatum* to enable gene expression study using real-time quantitative RT-PCR. *Curr Genet* 60(3):231–236
- Zhang JH, Shi JL, Liu YL (2013a) Bioconversion of resveratrol using resting cells of nongenetically modified *Alternaria* sp. *Biotechnol Appl Biochem* 60(2):236–243
- Zhang JH, Shi JL, Liu YL (2013b) Substrates and enzyme activities related to biotransformation of resveratrol from phenylalanine by *Alternaria* sp. MG1. *Appl Microbiol Biotechnol* 97(23):9941–9954
- Zhou YH, Zhang YJ, Luo ZB, Fan YH, Tang GR, Liu LJ, Pei Y (2012) Selection of optimal reference genes for expression analysis in the entomopathogenic fungus *Beauveria bassiana* during development, under changing nutrient conditions, and after exposure to abiotic stresses. *Appl Microbiol Biotechnol* 93(2):679–685

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)