

EVALUATION OF AN INDIRECT ELISA FOR THE DETECTION OF *SALMONELLA* IN CHICKEN MEAT

Andréa dos Santos Schneid¹; Kelly Lameiro Rodrigues¹; Davi Chemello²; Eduardo Cesar Tondo²;
Marco Antônio Zacchia Ayub²; José Antonio Guimarães Aleixo^{1*}

¹Centro de Biotecnologia, Universidade Federal de Pelotas, Pelotas, RS, Brasil; ²Instituto de Ciência e Tecnologia de Alimentos, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brasil

Submitted: March 15, 2005; Approved: June 14, 2006

ABSTRACT

In this work, an indirect ELISA based on a monoclonal antibody (MAb) specific for an outer membrane protein of *Salmonella enterica* serovar Enteritidis was used for detection of *Salmonella* in 154 samples of chicken meat. Its efficiency was determined through comparison with the results obtained from the conventional method. The prevalence of samples contaminated with *Salmonella* was 23% with the conventional culture method, and 26% with the ELISA. From thirty-five samples positive for *Salmonella* by the conventional method, 33 were also positive by ELISA. Seven other samples were only positive in the ELISA. Comparison of the results obtained in the two methods showed an ELISA sensitivity and specificity of 94%, and positive and negative predictive values of 82% and 98% respectively. The serotyping of the isolates revealed 31 *Salmonella enterica* serovar Enteritidis, 2 *Salmonella enterica* serovar Heidelberg, 1 *Salmonella enterica* serovar Choleraesuis and 1 *Salmonella enterica* serovar 6,7:-:-.

Key words: *Salmonella*, monoclonal antibody, policlonal antibody, ELISA

INTRODUCTION

Salmonella bacteria are often involved in outbreaks of gastroenteritis all over the world (21,28). As *Salmonella* is widely spread throughout the environment, contaminating water and a wide range of foods, it represents a serious problem for public health and it is therefore necessary to have fast and reliable methods of detection.

The conventional method of cultivation used in detection is reliable but slow as it includes stages of pre-enrichment, selective enrichment, cultivation in selective agars, biochemical characterisation of suspected isolates and a final serological confirmation (20). The relatively long time required to carry out analysis (4-7 days) as well as the costs incurred and the storage of foods suspected of being contaminated have stimulated the development of faster detection methods that can be useful in studies of the source of contamination and of outbreaks due to food contamination (7,15,19).

Among the methods that have already been put forward are DNA hybridisation (1,17), polymerase chain reaction (3,30,35), tests with specific bacteriophages (4), latex agglutination tests (9,11,32), immunomagnetic assays (29,34) and other types of immunoenzymatic assays (24,33,36,37).

The enzyme-linked immunosorbent assay (ELISA), a type of immunoenzymatic assay that uses solid phases to anchor the antibody-antigen complex, have been widely used because offer several advantages. One of these is the capacity to detect the presence of bacteria antigens without the cells having to be viable (24). Besides, many ELISA tests, mainly those that use polyclonal antibodies, show good sensitivity, but have restricted specificity for certain serovars. Others use a combination of polyclonal antibodies in order to obtain a wide spectrum of specificity. However, this method may potentialize the occurrence of unspecific reactions due to the nature and the way that these antibodies are obtained (6).

*Corresponding Author. Mailing address: Centro de Biotecnologia, Universidade Federal de Pelotas, Campus Universitário. 96030-001. Pelotas, RS, Brasil. Tel.: (+5553) 275-7350. Fax: (+5553) 275-7355. E-mail: biotjaga@ufpel.tche.br

The use of monoclonal antibodies (MAbs) with high specificity and affinity has been a good alternative in making ELISA tests more reliable. The use of a combination of MAbs for specific *Salmonella* epitopes is also an important tool in order to increase the sensitivity of the ELISA (10).

This study aims to evaluate an ELISA system that uses a specific monoclonal antibody for a protein taken from the external membrane of *Salmonella* Enteritidis in the detection of *Salmonella* in naturally contaminated chicken.

MATERIALS AND METHODS

Sample collection and preparation

The flow chart of procedures (20) used during the collection and processing of samples is shown in Fig. 1. A total of 154 samples of refrigerated chicken cuts were obtained from a local shop. The samples, thighs and legs, which are normally on display in trays on refrigerated shelves, were bought and

wrapped in plastic film and then placed in polystyrene with ice for transportation to the laboratory where the analyses were begun on the same day. The aseptic collection of 25g samples was carried out on a laminar flow bench. The samples were then placed in sterile plastic bags containing 225 mL of lactose broth (LB) (Difco Laboratories, Detroit, MI), homogenised for two minutes in Stomacher™, incubated at 37°C for 18-24 h and from there on the detection stages of the two methods being compared were done.

Conventional detection method

After the incubation period, 0.1 mL and 1.0 mL of cultures in LB (Difco) were transferred, respectively, into Rappaport Vassiliadis (RV) (Difco Laboratories, Detroit, MI) and tetrionate (TT) (Difco) broths and incubated at a temperature between 42°C (RV) and 43°C (TT) for 18 - 24 h. At the end of the period of selective enrichment, the broths were plated in Hektoen enteric agar (Difco) and bismuth sulphite agar (Difco) and incubated at 37°C for 18 - 24h in order to isolate suspected colonies. Biochemical identification was then carried out on 3 to 5 isolated colonies by cultivation at 37°C for 24 h in triple sugar iron agar (TSI) (Difco), lysine iron agar (LIA) (Difco) and urea broth (Difco). Serological confirmation of isolates was carried out by slide agglutination using somatic polyvalent serum and in tubes using polyvalent flagellar serum (Probac, São Paulo, SP). The final serotype identification was carried out at the Adolfo Lutz Institute (São Paulo, Brazil).

ELISA procedures

After selective enrichment, portions of 0.5 mL of RV and TT were transferred into tubes containing 10 mL of brain heart infusion broth (BHI) in order to carry out post-enrichment for 6h at 37°C in a water bath with agitation. The post-enriched cultures were heated (100°C/10 min) and used to sensitize four wells of polystyrene plates with 50 µL/well for 1h at 37°C. The plates were washed three times with phosphate-buffered saline (0.01 M, pH 7.2) containing 0.05% Tween 20 (PBST), and two wells received MAb 424H, that react specifically with a putative outer membrane protein from salmonellae (27), and the other two were used without MAb for control of unspecific reactions. The wells received 50 µL of the MAb diluted 1:1000 in PBST and left for 1h at 37°C. After washing the plates another three times with PBST, 50 µL/well of a protein A-peroxide conjugate (Sigma, Saint Louis, MO) diluted 1:2000 in PBST was added and left for 1h at 37°C. The plates were washed a further 5 times with PBST, and 50 µL well of a cromogene/substrate solution was added [40 mg of ortophenylenediamine dihydrochloride (OPD), 40 µL of hydrogen peroxide- H₂O₂ and 100 mL of citrate-phosphate buffer 0.05M, pH 4.5] and left to react for 15 min in the dark at room temperature. The optical density (OD) was then read at 450 nm in a MR 700 microplate reader (Dynatech Laboratories, INC., Virginia). The ELISA cut off for positive




Figure 1. Flow chart used to compare the detection of *Salmonella* in chicken meat by conventional methodology and indirect ELISA. RV: Rappaport Vassiliadis broth; TT: tetrionate broth; HE: Hektoen agar; BS: bismuth sulphite agar; TSI: triple sugar iron agar; LIA: lysine iron agar; BHI: brain heart infusion broth.

samples corresponded to a mean OD above 0.095, or the 96th percentile of 20 negative samples (27). As ELISA positive control, each plate contained four wells sensitized with a thermo-extracted antigen from *Salmonella* Enteritidis ATCC 13076. The control antigen was obtained by heating a suspension of cells with an OD₆₀₀ of 1.00, prepared from a culture in Brain Heart Infusion broth (BHI; Difco Laboratories, Detroit, MI) at 37°C for 6h. As ELISA negative control, four wells of a mixture of antigens heat-extracted from BHI cultures of chicken samples negative for salmonellae were used. The ELISA was considered to be acceptable when the mean OD of the positive control wells was between 1.0 and 1.2 and of the negative control wells was 0.095 or less.

Statistical analysis

The data were computed using the Epi Info programme version 6.04 (13). The Yates X₂ correction test with a confidence interval of 95% was used to see if there had been significant differences between the detection results of *Salmonella* in the samples of chicken meat using the ELISA and the conventional methodology. Sensitivity (s), specificity (sp), predictive positive value (ppv) and negative predictive value (npv) of the ELISA in the 154 samples tested were defined as: (s) probability of obtaining a positive result if the sample was truly positive, (sp) probability of obtaining a negative result if the sample was truly negative, (ppv) probability that an ELISA positive sample was truly positive, (npv) probability that an ELISA negative sample was truly negative. Accuracy, which is defined as the percentage of results that coincide in the two methods, was obtained using the rate between the number of coincidental results in the ELISA and the conventional method, and the total number of positive and negative results by the conventional method.

RESULTS

The results obtained from the analysis of the 154 chicken samples of are shown in Table 1. The prevalence of samples contaminated with *Salmonella* was 23% using the conventional method of cultivation and 26% using the indirect ELISA test. This difference was not statistically significant ($P < 0.001$). The ELISA gave 7 false positive results and only 2 false negative results. Among the 35 samples where *Salmonella* was detected, 31 were contaminated with *Salmonella* Enteritidis, 2 with *Salmonella* Heidelberg, 1 with *Salmonella* Choleraesuis and 1 with *Salmonella enterica* strain 6.7:-:-.

Compared to the results obtained by the conventional method, the ELISA accuracy was 94% with sensitivity and specificity also of 94%, and predictive positive and negative values of 82% and 98% respectively (Table 2).

Table 1. Detection of *Salmonella* in chicken meat samples by indirect ELISA and conventional methodology.

Samples ^a	ELISA (OD ₄₅₀)	Conventional Methodology	Serovar
1	0.477	+	<i>Salmonella</i> Enteritidis
2	0.340	+	<i>Salmonella</i> Enteritidis
3	0.347	+	<i>Salmonella</i> Enteritidis
4	0.170	+	<i>Salmonella</i> Enteritidis
5	0.143	+	<i>Salmonella</i> Enteritidis
6	0.361	+	<i>Salmonella</i> Enteritidis
7	0.298	+	<i>Salmonella</i> Enteritidis
8	0.788	+	<i>Salmonella</i> Enteritidis
9	0.306	+	<i>Salmonella</i> Enteritidis
10	0.255	+	<i>Salmonella</i> Enteritidis
11	0.246	+	<i>Salmonella</i> Enteritidis
12	0.223	+	<i>Salmonella</i> Enteritidis
13	0.123	+	<i>Salmonella</i> Enteritidis
14	0.186	+	<i>Salmonella</i> Enteritidis
15	0.183	+	<i>Salmonella</i> Enteritidis
16 ^b	0.063	+	<i>Salmonella</i> Enteritidis
17	0.421	+	<i>Salmonella</i> Enteritidis
18	0.420	+	<i>Salmonella</i> Enteritidis
19	0.275	+	<i>Salmonella</i> Enteritidis
20	0.270	+	<i>Salmonella</i> Enteritidis
21	0.107	+	<i>Salmonella</i> Enteritidis
22	0.757	+	<i>Salmonella</i> Enteritidis
23	0.286	+	<i>Salmonella</i> Enteritidis
24	0.252	+	<i>Salmonella</i> Enteritidis
25	0.242	+	<i>Salmonella</i> Enteritidis
26	0.265	+	<i>Salmonella</i> Enteritidis
27	0.342	+	<i>Salmonella</i> Enteritidis
28 ^b	0.090	+	<i>Salmonella</i> Enteritidis
29	0.533	+	<i>Salmonella</i> Enteritidis
30	0.334	+	<i>Salmonella</i> Enteritidis
31	0.576	+	<i>Salmonella</i> Enteritidis
32	0.782	+	<i>Salmonella</i> Choleraesuis
33	0.278	+	<i>Salmonella</i> Heidelberg
34	0.311	+	<i>Salmonella</i> Heidelberg
35	0.107	+	<i>Salmonella enterica</i> serovar 6,7:-:-
36 ^c	0.224	-	NI
37 ^c	0.109	-	NI
38 ^c	0.116	-	NI
39 ^c	0.263	-	NI
40 ^c	0.104	-	NI
41 ^c	0.107	-	NI
42 ^c	0.105	-	NI
112 Samples	< 0.095	-	-

OD₄₅₀ values over 0.095 are considered positives in ELISA; NI (no isolated); ^a A total of 154 samples of chicken meat were analysed; ^b false negative results in ELISA; ^c false positive results in ELISA.

Table 2. Sensitivity, specificity, positive and negative predictive values of indirect ELISA for detection of *Salmonella* in chicken meat samples.

Conventional ELISA	Methodology		
	positive	negative	total
Positive	33 ^a	7 ^b	40
Negative	2 ^c	112 ^d	114
Total	35	119	154

^apositive results in ELISA; ^bfalse positive results in ELISA; ^cfalse negative results in ELISA; ^dnegative results in ELISA; sensitivity ($a / a + c$) = 0.94; specificity ($d / b + d$) = 0.94; positive predictive value; ($a / a + b$) = 0.82; negative predictive value ($d / c + d$) = 0.98.

DISCUSSION

When comparing the results obtained by the ELISA and the conventional method in the analysis of the 154 chicken samples for the detection of *Salmonella*, one can see that they were not statistically different. According to these results, using the conventional method the prevalence of *Salmonella* in chicken was 23% while using the ELISA it was 26%. Both detection systems revealed a high percentage of chicken cuts contaminated with *Salmonella*. This finding proves how important this type of food probably is in the transmission of this bacteria to humans in our community. Animal based foods, especially chicken, have been associated with the occurrence of *Salmonella* in humans, *Salmonella* Enteritidis being the most frequently isolated strain in this type of food (8). Similar results were found in this study which showed that out of a total of 35 *Salmonella* positive samples, 31 were of the *Salmonella* Enteritidis strain which in turn corresponds to 88.6% of the strains isolated.

The results reveal that the ELISA and the conventional methodology behaved in a similar fashion in terms of efficiency. The ELISA was able to detect *Salmonella* in 40 samples, 7 more than the conventional system, a number of false positive results reasonably low (4.5%). This is reflected in the high values obtained for specificity (94%) and the positive prediction value (82%) (Table 2). Various studies comparing ELISA and conventional testing verify a tendency for ELISA tests to give false positive results (4,7,25). This tendency is generally attributed to cross reactions of antibodies with antigens of others microorganisms present in the samples or to reactions with constituents in the food. However, the low number of *Salmonella* in the samples, cellular injury and the interference of natural flora in the multiplication of *Salmonella* in foods, are some factors that may contribute to difficulties or prevention of isolation thereby causing false negative results in the conventional method (14,23). Also, the isolation of pathogens

in some agars may be made difficult by the presence of a large number of competitive bacteria, some of which are very similar to *Salmonella* and which may cause error when selecting suspect colonies (26,31).

Peplow *et al.* (25) stated that false negative results in rapid detection methods are easily proved by isolation of microorganisms using the conventional method and that, on the contrary, false positive results in rapid detection methods cannot be proved as they are, in reality, false negatives in the conventional methodology. It is therefore possible that the false positive results observed in our ELISA may be due to a detection fault in the conventional test. According to Bailey (2), the new technologies for detection of pathogens in foods are more sensitive than conventional methods besides eliminating the need for isolation to confirm the presence of pathogens.

The two false negative results given by the ELISA test (1.3%) directly influenced sensitivity (94%) and the predictive negative value (98%). One explanation for the false negatives may have been the low number of *Salmonella* cells present in the post enrichment phase. It is possible that the natural flora present in these two samples impaired the multiplication of *Salmonella* cells at levels that were detectable by ELISA (16). It is also possible that the MAb 424H in the reaction had not retained enough enzyme to form a sufficient quantity of product to be detected by the system. An alternative to avoid this type of error is to use a combination of antibodies that are specific for *Salmonella* antigens in order to amplify test results by the higher number of epitopes detected (5). Another alternative would be the improvement of enrichment protocols used in the ELISA test by inhibiting microbial competition and favouring the growth of *Salmonella* to increase the amount of antigen available for reaction with detector antibody (12,18,22,31).

In conclusion, the indirect ELISA with the MAb 424H used to detect the presence of *Salmonella* in chicken cuts allows positive results 30 hours before conventional testing. The ELISA has a sensitivity and specificity comparable to other ELISA tests already developed and commercially available. It has a good potential for use as a screening test in the quality control of the food industry.

RESUMO

Avaliação de um ELISA indireto para detecção de *Salmonella* em carne de frango

Neste trabalho, um ELISA indireto baseado em um anticorpo monoclonal (MAB) específico para proteína de membrana externa de *Salmonella enterica* serovar Enteritidis foi usado para detecção de *Salmonella* em 154 amostras de carne de frango. Sua eficiência foi determinada através de comparação com os resultados obtidos pela metodologia convencional. A

prevalência de amostras contaminadas com *Salmonella* foi de 23% pelo método de cultivo convencional, e 26% pelo ELISA. De 35 amostras positivas para *Salmonella* pela metodologia convencional, 32 também foram positivas no ELISA. Outras sete amostras foram positivas somente no ELISA. Comparando os resultados obtidos nos dois métodos, o ELISA demonstrou sensibilidade e especificidade de 94%, e valor preditivo positivo e negativo de 82% e 98% respectivamente. A sorotipagem dos isolados revelou 31 *Salmonella enterica* serovar Enteritidis, 2 *Salmonella enterica* serovar Heidelberg, 1 *Salmonella enterica* serovar Choleraesuis e 1 *Salmonella enterica* serovar 6,7:-:-.

Palavras-chave: *Salmonella*, anticorpo monoclonal, anticorpo policlonal, ELISA

REFERENCES

- Bailey, J.S.; Cox, N.A.; Blankenship, L.C. A comparison of an enzyme Immunoassay, DNA hybridization, antibody immobilization, and conventional methods for recovery of naturally occurring *Salmonella* from processed broiler carcasses. *J. Food Protect.*, 54, 354-356, 1991.
- Bailey, J.S. Detection of *Salmonella* cells within 24 to 26 hours in poultry samples with the polymerase chain reaction BAX system. *J. Food Protect.*, 61(7), 792-795, 1998.
- Baumler, A.J.; Heffron, F.; Reissbrodt, R. Rapid detection of *Salmonella enterica* with primers specific for iroB. *J. Clin. Microbiol.*, 35(5), 1224-1230, 1997.
- Bennett, A.R.; Davids, F.G.C.; Vlahodimou, S.; Banks, J.G.; Betts, R.P. The use of bacteriophage-based systems for the separation and concentration of *Salmonella*. *J. Appl. Bacteriol.*, 83, 259-265, 1997.
- Blackburn, C.W. A review. Rapid and alternative methods for the detection of salmonelas in foods. *J. Appl. Bacteriol.*, 75, 199-214, 1993.
- Brown, J.G.; Brooks, B.W.; Blais, B.W.; Yamazaki, H. Application of Cloth- based enzyme immunoassay for the characterization of monoclonal antibodies to *Salmonella* lipopolysaccharide antigens. *Immunol. Investigations*, 25(4), 369-381, 1996.
- Chaicumpa, W.; Ngren-Ngarmlert, W.; Kalambaheti, T.; Chongsanguan, Y.R.M.; Tapchaisri, P.; Desakorn, V.; Suthienkul, O. Monoclonal antibody - based Dot-Blot ELISA for the detection of *Salmonella* in foods. *Asian Pacific J. Allergy Immunol.*, 13, 159-166, 1995.
- Chang, Y.H. Prevalence of *Salmonella* spp. in poultry broilers and shell eggs in Korea. *J. Food Protect.*, 63(5), 655-658, 2000.
- Cheesbrough, S.; Donnelly, C. The use of a rapid *Salmonella* latex serogrouping test (Spectate) to assist in the confirmation of ELISA-based rapid *Salmonella* screenig tests. *Lett. Appl. Microbiol.*, 22(5), 378-380, 1996.
- Choi, D.; Tsang, R.S.W.; NG, M.H. Sandwich capture ELISA by a murine monoclonal antibody against a genus-specific LPS epitope fo the detection of different common serotypes of salmonellas. *J. Appl. Bacteriol.*, 72, 134-138, 1992.
- D'Aoust, J.Y.; Sewell, A.M.; Greco, P. Commercial latex agglutination kits for the detection of foodborne *Salmonella*. *J. Food Protect.*, 54(9), 725-730, 1991.
- D'Aoust, J.Y.; Sewell, A.N.; Daley, E. Inadequacy of small transfer volume and short (6h) selective enrichment for the detection of foodborne *Salmonella*. *J. Food Protect.*, 55(5), 326-328, 1992.
- Deam, A.G.; Deam, J.A.; Coulmobier, D.; Brendel, K.A.; Smith, D.C.; Burton, A.H.; Dicker, R.C.; Sullivan, K.; Fagan, R.F.; Arner, T.J. Epi Info version 6.0. A word processing, database, and statistics program for epidemiology on microcomputers. Centers for Disease Control and Prevention, Atlanta, GA, 1994.
- Eckner, K.F.; Dustman, W.A.; Curiale, M.S.; Flowers, R.S.; Robison, B.J. Elevated-temperature, colorimetric, monoclonal, enzyme-linked immunosorbent assay for rapid screening of *Salmonella* in foods: collaborative study. *J. Assoc. Off. Anal. Chem.*, 77(2), 374-383, 1994.
- Fach, P.; Dilasser, F.; Grout, J.; Tache, J. Evaluation of a polimerase chain reaction-based test for detecting *Salmonella* spp in foods samples: Probelia *Salmonella* spp. *J. Food Protect.*, 62(12), 1387-1393, 1999.
- Feldsine, P.T.; Nelson, M.T.F.; Husted, D.L. Policlonal enzyme immunoassay method for detection of motile and non-motile *Salmonella* in foods: collaborative study. *J. Assoc. Off. Anal. Chem.*, 76(4), 694-697, 1993.
- Fitts, R.; Diamond, M.; Hamilton, C.; Neri, M. DNA - DNA Hibridization assay for detection of *Salmonella* spp. in foods. *Appl. Environment. Microbiol.*, 46, 1146-1149, 1983.
- Flowers, R.S.; Klatt, M.J.; Robison, B.J.; Mattingly, J.A. Evaluation of abbreviated enzyme immunoassay method for detection of *Salmonella* in low-moisture foods. *J. Assoc. Off. Anal. Chem.*, 71(4), 341-343, 1988.
- Flowers, R.S.; Klatt, M.J.; Keelan, S.L.; Gehle, W.D.; Chandonnet, E. Fluorescent enzyme immunoassay for rapid screenin of *Salmonella* in foods: Collaborative study. *J. Assoc. Off. Anal. Chem.*, 72(2), 318-325, 1989.
- Food and Drug Administration. Bacteriological Analytical Manual, 8th Ed. AOAC International, Arlington, 1998.
- Guo, X.; Chen, J.; Beuchat, L.R.; Brackett, R.E. PCR detection of *Salmonella enterica* serotipo Montevideo in and on raw tomatoes using primers derived from *hlyA*. *Appl. Environ. Microbiol.*, 66(12), 5248-5252, 2000.
- Kerr, S.; Ball, H.J.; Mackie, D.P.; Pollock, D.A.; Finlay, D.A. Diagnostic application of monoclonal antibodies to outer membrane protein for rapid detection of *salmonella*. *J. Appl. Bacteriol.*, 72, 302-308, 1992.
- Krusell, L.; Skovgaard, N. Evaluation of a new semi-automated screening methods for the detection of *Salmonella* in foods within 24 h. *Int. J. Food Microbiol.*, 20, 123-130, 1993.
- Ng, S.P.; Tsui, C.O.; Roberts, D.; Chau, P.Y.; Ng, M.H. Detection and serogroup differentiation of *Salmonella* spp. in food within 30 hours by enrichment-immunoassay with a T6 monoclonal antibody capture Enzyme-Linked Immunosorbent Assay. *Appl. Environ. Microbiol.*, 62(7), 2294-2302, 1996.
- Peplow, M.O.; Prisant, M.C.; Stebbins, M.E.; Jones, F.; Davies, P. Sensitive, specificity, and predictive values of three *Salmonella* rapid detection kits using fresh and frozen poultry environmental samples versus those of standard plating. *Appl. Environ. Microbiol.*, 65(3), 1055-1060, 1999.
- Rambach, A. New plate medium for facilitated differentiation of *Salmonella* spp. from *Proteus* spp. and other enteric bacteria. *Appl. Environ. Microbiol.*, 56, 301-303, 1990.
- Schneid, A.S.; Lütke, C.B.; Diel, C.; Aleixo, J.A.G. Production and characterization of monoclonal antibodies for the detection of *Salmonella enterica* in chicken meat. *Braz. J. Microbiol.*, 36, 163-169, 2005.
- Seo, K.H.; Hartman, N.F.; Campbell, D.P. Development of a rapid response biosensor for detection of *Salmonella* Typhimurium. *J. Food Protect.*, 62(5), 431-437, 1999.
- Shaw, J.S.; Blais, B.W.; Nundy, D.C. Performance of the Dynabeads anti-*Salmonella* systems in the detection of *Salmonella* species in foods, animal feeds, and environmental samples. *J. Food Protect.*, 61(11), 1507-1510, 1998.

30. Stone, G.G.; Oberst, R.D.; Hays, M.P.; McVey, S.; Chengappa, M.M. Combined PCR-oligonucleotide ligation assay for rapid detection of *Salmonella* serovars. *J. Clin. Microbiol.*, 33(11), 2888-2893, 1995.
31. Tate, C.R.; Miller, R.G.; Mallinson, E.T. Evaluation of two isolation and two no-isolation methods for detecting naturally occurring *Salmonellae* from broiler flock environmental drag-swab samples. *J. Food Protect.*, 55, 964-967, 1992.
32. Thorns, C.J.; McLaren, I.M.; Sojka, M.G. The use of latex particle agglutination to specifically detect *Salmonella enteritidis*. *Int. J. Food Microbiol.*, 21, 47-53, 1994.
33. Veling, J.; Van Zijderveld, F.G.; Van Zijderveld-Van Bommel, A.M.; Barkema, H.W.; Chukken, Y.H. Evaluation of three Newly developed Enzyme-Linked Immunosorbent Assay and two agglutination tests for detecting *Salmonella enterica* subsp. *enterica* serovar Dublin infections in Dairy cattle. *J. Clin. Microbiol.*, 38(12), 4402-4407, 2000.
34. Vermunt, A.E.M.; Franken, A.A.J.M.; Beumer, R.R. Isolation of salmonelas by immunomagnetic separation. *J. Appl. Bacteriol.*, 72, 112-118, 1992.
35. Way, J.S.; Josephson, K.L.; Pillai, S.D.; Abbaszadegan, M.; Gerba, C.P.; Pepper, I.L. Specific detection of *Salmonella* spp by multiplex polymerase chain reaction. *Appl. Environ. Microbiol.*, 59(5), 1473-1479, 1993.
36. Wyatt, G.M.; Langley, M.N.; Lee, H.A.; Morgan, M.R.A. Further studies on the feasibility of one-day *Salmonella* detection by enzyme-linked immunosorbent assay. *Appl. Environ. Microbiol.*, 59, 83-1390, 1993.
37. Wyatt, G.M.; Lee, H.A.; Dionysiou, S.; Morgan, M.R.A.; Stokely, D.J.; Al-Hajji, A.H.; Richards, J.; Sillis, A.J.; Jones, P.H. Comparison of a microtitration plate ELISA with a standard cultural procedure for the detection of *Salmonella* spp. in chicken. *J. Food Protect.*, 59(3), 238-243, 1996.